

# Temporal Investigation of the Frequency-magnitude Distribution and Fractal Dimension in the Northern Thailand: A comparison

Kanokkarn Vejchakom\* and Santi Pailoplee

*Morphology of Earth Surface and Advanced Geohazards in Southeast Asia Research Unit (MESA RU), Department of Geology, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand*

\* Corresponding author E-mail: k.vejchakom@gmail.com

## ABSTRACT

According to the Indo-Australian and Eurasian plate collision, a large number of seismogenic fault zone were dominant as the intraplate earthquake sources within the Mainland Southeast Asia or the Thailand-Laos-Myanmar border. Up to the present, hazardous earthquakes were generated continuously in 35 yr during 1980-2015. Therefore, this study aims mainly to investigate the earthquake activates and pattern of earthquake occurrence in this region. Regarding to statistical seismology, both  $b$  value of the frequency-magnitude distribution and fractal dimension ( $D_c$ ) were, therefore, analyzed temporally. In addition in terms of the utilized dataset, both dataset of before- and after-declustering were recognized. As a result, it reveals that both  $b$  and  $D_c$  value analyzed by the after-declustering dataset is meaningless in term of seismotectonic meaning. In contrast, the before-declustering dataset is meaningful for both  $b$  and  $D_c$  analysis. Based on the before-declustering dataset,  $b$  value is decrease continuously during 35 yr of 1980-2015 before both Mae Lao and Tarlay earthquakes posed. In addition, the  $D_c$  values are drop down suddenly during or after these earthquakes were generated. This may imply the earthquake precursor which useful for earthquake mitigation in any site of interest the future. For  $b$ - $D_c$  relationships, both Mae Lao and Tarlay earthquakes express the different correlation. Mae Lao earthquake indicate the positive relation while Tarlay earthquake express negative relation. This relation, therefore, may accord to the different characteristic or activities of each fault zone. To further refine seismic activities ( $b$  value) and seismic pattern ( $D_c$ ) value including their relationship in this region, more detailed case studies of earthquake that generated in different seismogenic fault are need.

**Keywords:** Earthquake Catalogue; Earthquake Declustering;  $b$  value; Fractal Dimension; Thailand-Myanmar Borders.

## 1. Introduction


The Indo-Australian plate has been moved and subducted northward underneath the Eurasian plate since 40 million years ago (Molnar and Topponnier, 1977). Seismotectonically, this plate collision initiates interplate earthquake sources called the Sumatra-Andaman Subduction Zone (Figure 1a). In addition, it also causes

intraplate earthquake sources where the inland seismogenic faults are dominant (Figure 1b). As an example of intraplate fault zones in the northern Thailand, there are the Mae Tha fault zones and also the Lampang-Thoen fault zone (Pailoplee et al., 2009). Therefore, a large number of earthquakes are generated continuously

within the Eurasian plate (Pailoplee et al., 2013).

According to previous research work, the TLMB which encompasses major cities and hydropower dams along the Mekong River (Mekong River Commission, 2010). There had generated a large number of hazardous earthquakes (Figure 1b) such

as magnitude 6.2 (1989), magnitude 6.8 (1995), magnitude 6.3 (2007), magnitude 6.8 (2011) (Pailoplee et al., 2013) and latest event is Mea Lao earthquake with magnitude 6.2 (2014) as shown in Figure 1b. The occurrence of earthquakes on the TLMB causes damage to northern Thailand, northern Laos and eastern Myanmar.



**Figure 1.** (a) Map of the mainland Southeast Asia showing the study area (blue square) and the boundary of earthquake investigation (black dash line square). (b) Map of study area showing the earthquake with  $M_w \geq 6.0$  (red circles) within the TLMB. The fault lines, hydro power dam, major cities are shown with thin grey lines, black triangles and black squares, respectively. The blue and green dots are the earthquake dataset occurred during 1965 to 2016 that before and after declustering process, respectively.

However in the study of earthquake characteristic or pattern, it is difficult to investigate the earthquake source in the TLMB because of the complexity in the seismotectonic setting and the blind faults made by the covering of quaternary sediment within the basin (Fenton et al., 2003). Scientists applied statistics to study earthquakes characteristics and estimate the changes of earthquakes activities in study area. In order to do so, they used methods such as the frequency-magnitude

distribution model (FMD; Nuannin et al., 2005), fractal dimension ( $D_c$ ; Chen et al., 2006), Pattern informatics (PI; Wu et al., 2008), seismic rate change ( $Z$ ; Wyss and Habermann, 1988), and region-time-length algorithm (RTL; Chen and Wu, 2006) etc. In addition, some previous works reported that the spatial distribution of  $b$  value and  $D_c$  values have a relationships implying the accumulated tectonic stress and seismic pattern, respectively (Pailoplee et al., 2014). The main aim of this study is, therefore,

analyzing the temporal variation of both of  $b$  and  $D_c$  values in the study area. The epicenters of 2 case studies of significant earthquakes were recognized, i.e., Mea Lao (2014) and Tarlay (2011) earthquakes. In addition in technical term, this study also considered both  $b$  and  $D_c$  based on 2 different earthquake dataset, i.e., dataset i) before and ii) after declustering.

## 2. Dataset and Completeness

### 2.1. Earthquake catalog collection

The earthquake catalogues are one of the most important products of seismology (Woessner and Wiemer, 2005) and they are important in the study of earthquake characteristics using statistical analysis. In this study, we used the earthquakes that occurred on the TLMB recorded from 1966 to 2016. This yielded approximately 12,133 recorded earthquake events (blue dots in Figure 1b). These catalogue were reported by i) the International Seismological Centre (ISC), ii) the National Earthquake Information Center (NEIC) and iii) the Global Centroid Moment Tensor (GCMT). The earthquake magnitude were reported variously in body-wave magnitude ( $M_b$ ), surface-wave magnitude ( $M_s$ ), local magnitude ( $M_l$ ) and moment magnitude ( $M_w$ ) scales. In order to homogenize the magnitude scale, all different scales were converted empirically to the  $M_w$  which represent directly the seismotectonic activities.

### 2.2. Earthquake decluttering

Normally, the earthquakes occurrences can be sorted out into 3 types which are i) foreshock, ii) mainshock and iii) aftershock. Seismically, only mainshocks indicate directly about the tectonic stress accumulated in any specific area (Nuannin et al., 2005). Therefore, both foreshock and aftershock clusters should be screen and remove from the utilized earthquake catalogue in any seismicity investigation in the statistical approach. According to the assumption of Gardner and Knopoff (1974), the earthquake dataset of

the TLMB were declustered statistically (Figure 2). As a result, the mainshock catalogue comprises a total of 2,195 earthquake events with  $M_w \geq 0.1$  from 1964 to 2016 (green dot in Figure 1b).

As mentioned above, both earthquake dataset of i) before and ii) after declustering were utilized in this analysis in order to test technically the possibility of using the dataset. Therefore, in this temporal investigation of the seismicity, i.e.,  $b$  and  $D_c$  values, the earthquake data within 300-km radius (Kupta, 2002) from the hazardous Mea Lao and Tarlay earthquakes were selected from both the earthquake data of the before- and after-declustering process. As a result, 4 case studies of temporal investigation were demonstrated in this analysis.


## 3. Frequency-magnitude Distribution ( $b$ Value)

According to Gutenberg and Richter (1944), the empirical relationship of the FMD is shown in Equation (1)



$$\log(N) = a - bM \quad (1)$$

Where  $N$  is the cumulative number of earthquakes with magnitude greater or equal to  $M$ .  $a$  and  $b$  are constants, positive and vary in any specific time and space window. The  $a$  value is all of the activity level of seismicity, and the  $b$  value is the slope of FMD. In seismological context,  $b$  value related to the variation of the tectonic stress accumulated (Mogi, 1967; Scholz, 1986). Lower  $b$  value relates with large tectonic stress, strain and faults (Manakou and Tsapanos, 2000).

For example in case of the after-declustering dataset within 300-km radius from the Mea Lao earthquake, the 492 earthquake data were recorded during 1965-2016. The FMD plot (Figure 3a) represented the y-intercept as a value is 3.13. The  $b$  value indicated by slope is  $0.634 \pm 0.04$ . The magnitude of completeness is 4.0.



**Figure 2.** Diagram showing the declustering process of the earthquake data in the TLMB according to Gardner and Knopoff’s declustering model (red line). The earthquake data (blue dots) upper both red lines of (a) different of occurrence time and (b) distance from each earthquake event were defined as the earthquake mainshocks. Meanwhile, the other data below the redline were classified as foreshocks or aftershocks.



**Figure 3.** (a) The FMD plot of the Mea lao after declustering dataset. The triangles and squares represent the number and cumulative number of each individual seismicity magnitude level, respectively. The red line indicates the FMD linear regression fitted with the observed data. The  $M_c$  is defined as the magnitude of completeness (Woessner and Wiemer, 2005). (b) Graph showing the relation between  $\log(C_r)$  and  $\log(R)$  of the Mea lao after declustering dataset. The slope of linear fit (green line) is the fractal dimension ( $D_c$ ).

**4. Fractal Dimension ( $D_c$  Value)**

The fractal dimension ( $D_c$  value) is estimated using the correlation dimension

by the correlation integral technique. This technique measures the spacing of a set of points are earthquake epicenters

(Grassberger and Procaccia, 1983). The correlation dimension is widely applied in spatial term of earthquake epicenters distribution. That is preferable because of its greater reliability and sensitive to small changes in clustering properties (Kagan and Knopoff, 1980; Hirata, 1989). So, the fractal dimension of epicenter distribution of earthquake was calculated from the correlation technique given relation by Grassberger and Procaccia (1983) is defined as Equation (2).

$$D = \frac{\lim_{r \rightarrow 0} \log(C_r)}{\log r} \quad (2)$$

Where  $(C_r)$  is the correlation function as expressed in Equation (3).

$$C_r = \frac{2(N_{(R < r)})}{N(N - 1)} \quad (3)$$

Where  $N_{(R < r)}$  is the number of points separate by distance  $R$  less than  $r$ .  $N$  is the total number of points analyzed. If the epicenters distribution of earthquake has a fractal structure, the correlation integral is related to the standard correlation function as illustrated in Equation (4).

$$C_r \sim r^{D_c} \quad (4)$$

Where  $D_c$  is a fractal dimension, more strictly, the correlation dimension (Grassberger and Procaccia, 1983). By plotting  $C_r$  against  $r$  on a double logarithmic

## 5. Result

### 5.1. Temporal variation of $b$ and $D_c$ values

As mentioned above, we focus on the temporal variation of the tectonic stress (i.e.,  $b$  value) and the seismic pattern ( $D_c$ ) before both Tarlay and Mea-Lao earthquakes. The earthquakes data was divided into temporal term and analysed both  $b$  and  $D_c$  as shown in Table 1

For the Mea Lao earthquake, the  $b$  values analyzed from the before-

coordinate, we can calculate practical fractal dimension value ( $D_c$ ) from slope of this graph.

The distance ( $r$ ) between two events,  $(\theta_1, \phi_1)$  and  $(\theta_2, \phi_2)$  are the colatitudes ( $\theta$ ) and longitudes ( $\phi$ ), is calculated by using the formula given by Hirata (1989) as expressed in Equation (5).

$$r = \cos^{-1}(\cos \theta_1 \cos \theta_2 + \sin \theta_1 \sin \theta_2 \cos(\phi_1 - \phi_2)) \quad (5)$$

The slope is obtained by fitting a least-square line in the scaling region. The fractal dimension  $D_c$  gives an estimate of the fractal characteristics of fault system. Seismically, the  $D_c$  values close to 3 designate the earthquake fractures are generated within the volume source of crust. Meanwhile, the  $D_c$  values close to 2 imply the earthquake occurrence is being filled up by plane. And also, the  $D_c$  values close to 1 means line sources (Aki, 1981).

For example in case of the after-declustering dataset within 300-km radius from the Mea Lao earthquake, i.e., the same dataset used in FMD plot in Figure 3a. We found the slope from the relation between distance ( $R$ ) and correlation integral ( $C_r$ ) as  $2.00 \pm 0.01$ . That slope indicated the fractal value ( $D$ ), considered in distance between 12.99 km. to 93.01 km as shown in Figure 3b. This  $D$  value specify the earthquake distribution pattern as plate source.

declustering dataset reveal that  $b$  value varies temporally during 1980-2016 with the  $b$  values are in the range of 0.62-1.40 (Table 1a and Figure 4a). The highest  $b$  value of 1.40 expresses at 1980. After that the  $b$  value decrease continuously until 2000. The  $b$  values are stable at 0.62-0.75 from 2000 to 2016 (Figure 4a). In case of the Tarlay earthquake, we found the highest  $b$  value ( $b = 1.08$ ) analyzed from the before-declustering is at 1985 (Table 1c and Figure 4c). After that,  $b$  values between 1990-2016 are dropped down to 0.65-0.70 (Figure 4c).

**Table 1.** The result of temporal  $M_c$ ,  $a$ ,  $b$ , and  $D_c$  investigation analyzed at the epicenters of the Mea Lao and Tarlay earthquakes using the earthquake dataset that before and after declustering process.

**(a) Mea Lao before declustering**

Year	EQ number	$M_c$	$a$	$b$	$D_c$
1965-1980	57	4.8	7.06	$1.40 \pm 0.20$	$0.65 \pm 0.01$
1965-1985	143	4.9	6.21	$1.20 \pm 0.10$	$1.45 \pm \text{NaN}$
1965-1990	326	4.6	4.73	$0.87 \pm 0.06$	$1.73 \pm \text{NaN}$
1965-1995	536	4.5	4.31	$0.75 \pm 0.04$	$1.38 \pm \text{NaN}$
1965-2000	682	4.0	3.68	$0.62 \pm 0.02$	$1.32 \pm 0.03$
1965-2005	858	4.2	4.23	$0.74 \pm 0.03$	$1.82 \pm 0.03$
1965-2010	1446	4.0	3.91	$0.66 \pm 0.02$	$1.75 \pm 0.03$
1965-2015	2516	4.0	3.92	$0.65 \pm 0.02$	$1.17 \pm 0.02$
1965-2016	2521	4.0	3.92	$0.65 \pm 0.02$	$1.55 \pm 0.07$

**(b) Mea Lao after declustering**

Year	EQ number	$M_c$	$a$	$b$	$D_c$
1965-2000	169	4.2	3.12	$0.63 \pm 0.05$	$2.01 \pm 0.02$
1965-2005	228	4.2	3.4	$0.69 \pm 0.05$	$2.00 \pm 0.02$
1965-2010	386	4.0	3.12	$0.63 \pm 0.04$	$1.98 \pm \text{NaN}$
1965-2015	490	4.0	3.12	$0.63 \pm 0.04$	$2.00 \pm 0.01$
1965-2016	492	4.0	3.13	$0.63 \pm 0.04$	$2.00 \pm 0.01$

**(c) Tarlay before declustering**

Year	EQ number	$M_c$	$a$	$b$	$D_c$
1965-1985	241	4.9	6.01	$1.08 \pm 0.09$	$1.12 \pm 0.01$
1965-1990	637	4.3	4.07	$0.65 \pm 0.02$	$1.89 \pm 0.02$
1965-1995	950	4.3	4.12	$0.65 \pm 0.02$	$1.87 \pm 0.02$
1965-2000	1168	4.3	4.21	$0.67 \pm 0.02$	$1.91 \pm 0.01$
1965-2005	1418	4.3	4.33	$0.70 \pm 0.02$	$1.92 \pm 0.03$
1965-2010	2262	4.3	4.35	$0.70 \pm 0.02$	$1.97 \pm 0.02$
1965-2015	3410	4.3	4.31	$0.68 \pm 0.02$	$1.35 \pm 0.03$
1965-2016	3420	4.3	4.33	$0.68 \pm 0.02$	$1.35 \pm 0.03$

**(d) Tarlay after declustering**


Year	EQ number	$M_c$	$a$	$b$	$D_c$
1965-1995	184	4.2	3.14	$0.59 \pm 0.04$	$1.91 \pm 0.01$
1965-2000	251	4.2	3.31	$0.63 \pm 0.04$	$2.10 \pm 0.01$
1965-2005	347	4.2	3.48	$0.67 \pm 0.04$	$1.99 \pm 0.03$
1965-2010	537	4.2	3.51	$0.67 \pm 0.04$	$2.05 \pm \text{NaN}$
1965-2015	640	4.0	3.23	$0.62 \pm 0.03$	$2.13 \pm 0.01$
1965-2016	643	4.0	3.23	$0.62 \pm 0.03$	$2.17 \pm 0.01$

For  $D_c$  investigation in the Mae Lao earthquake location, the  $D_c$  values varies between 0.65-1.75. At 1980,  $D_c$  is 0.65 and increase continuously to 1990 at  $D_c = 1.73$

(Table 1a and Figure 4a). Thereafter  $D_c$  decrease during 1995-2000 down to 1.32-1.38. and increase up to 1.75-1.82 at 2005-2010 (Figure 4a). However during and after

the Mae Lao earthquake was generated at 2014, the Dc value is decrease again down to 1.17-1.15 (Figure 4a). Similar to the Dc values estimated at the Talay earthquake (Table 1c and Figure 4c), the Dc varies between 1.12-1.97. At the short time of 1965-1985, the Dc is comparatively low at 1.12. After that, 20 yr during 1990-2010, the Dc values reveal the comparatively high at Dc = 1.98-2.01 (Figure 4b). In final, the Dc value drop down to 2.13-2.17 before the Talay earthquake was generated.

Regarding to the after-declustering dataset, both b and Dc value analyzed here are insufficient variation. All results show the stable b and Dc in both Mae Lao and Talay case studies as illustrated in Figures 4b and d (see also Table 1b and d). At the Mae Lao earthquake epicenter, b value is constant around 0.63-0.69 while Dc value is around 1.98-2.01 (Figures 4b). For Talay earthquake, b value is at 0.59-0.67. Although Dc value is in the range of 1.91-2.17, this Dc range do not show any significant variation (Figure 4d).



**Figure 4.** Graphs showing the temporal variation of b (black line) and Dc (grey line) values from 4 case studies. The utilized datasets are comprise of (a) Mea Lao before declustering, (b) Mea Lao after declustering, (c) Talay before declustering, and (d) Talay after declustering.

### 5.2. b and Dc relationship


The b-Dc relationship has been suggested as an effective indicator of seismic hazards (Bayrak and Bayrak, 2011;

2012). Empirically, the b-Dc relationship can be either a positive or a negative correlation. For example, a positive correlation was defined at the earthquake

source zone in Northeast India (Bhattacharya et al., 2010). A negative correlation was revealed at some seismogenic source in Japan (Hirata, 1989) and the volcanic earthquake at Long Valley Caldera in California, USA (Barton et al., 1999).

In this study, we obtained the relations between  $b$  and  $D_c$  values of the Mea Lao and Tarlay earthquake sources (Figure 5). In case of Mae Lao earthquake, the relationship between  $b$  and  $D_c$  of the before-declustering dataset is  $D_c = 1.42b + 0.53$  (Figures 5a).

the after-declustering dataset, the  $b$ - $D_c$  relationship is  $D_c = 0.14b + 1.91$  (Figures 5b). These relationships are contributed based on the  $R^2$  0.24 and 0.16, respectively. Seismotectonically, the  $b$ - $D_c$  relationship of the Mae Lao earthquake is the positive relation indicating that low/high  $b$  value relates to low/high  $D_c$  value (Figures 5a and b). The positive correlation of Mae Lao earthquake conforms to the previous work that investigated at the earthquake source zone in Northeast India (Bhattacharya et al., 2010)



**Figure 5.** Empirical relationships between the  $b$  and  $D_c$  values analyzed from (a) Mea Lao before declustering, (b) Mea Lao after declustering, (c) Tarlay before declustering, and (d) Tarlay after declustering. The straight lines represent the linear regressions fitted with the observed data.

Regarding to the epicenter of the Tarlay earthquake, the relationship of the before-declustering dataset is  $D_c = -1.55b +$

$2.80$  with  $R^2 = 0.43$ . Whereas the after-declustering dataset reveals the  $b$ - $D_c$  relationship as  $D_c = -2.33b + 3.59$  with  $R^2$



= 0.81. In contrast to the Mae Lao earthquake, the b-Dc relationship of the Tarlay earthquakes expresses the negative linear correlation (Figures 5c and d). The negative correlation of Tarlay earthquake conforms to the previous works that reported the negative correlation at the region of seismogenic fault in Japan (Hirata, 1989) and the volcanic earthquake at Long Valley Caldera in California, USA (Barton et al., 1999).

## 6. Conclusion

In this study, the parameters of FMD (a and b values) and the fractal dimension (Dc value) were analyzed temporally at the epicenters of 2 well-known earthquakes, i.e., Mae Lao and Tarlay earthquakes. In addition in order to check the possibility of the utilized dataset, 2 earthquake datasets were employed, i.e., before- and after-declustering datasets. Based on the before-declustering dataset, the temporal variation of b value shows fairly implication. Before both Mae Lao and Tarlay earthquakes were generated, b values were decreased continuously throughout 30 yr during 1985-2015 (Figures 4a and c). It implies that the tectonic stress was accumulated within 300-km radius from each earthquake epicenter. Meanwhile for Dc value, there are some fluctuations of the Dc value during 1985-2015. And also, there is a significant drop of Dc value noticeable during the short time period (i.e., 5-10 yr) before these earthquakes were generated (Figures 4a and c). In contrast to the investigation using the before-declustering dataset, both b and Dc values analyzed from the after-declustering dataset do not show any meaningful variation of b and Dc in temporal. Consequently, we concluded that the before-declustering dataset can be used to analyze temporally both b and Dc values that imply the earthquake precursor prior to any hazardous earthquake posed.

Seismotectonically, b-Dc relationships reveal fairly correlation. However, each earthquake case study shows different relations. Mae Lao earthquake

indicates the positive relation while Tarlay earthquake expresses negative relation. Based mainly on the existing of 2 case studies, we conclude preliminarily that although the hazardous earthquakes were generated in the same seismotectonic setting (i.e., intraplate earthquake source) the relation between b and Dc values may differ. This may be according to the different characteristics or activities of each fault zone. The Mae Lao earthquake generated by Phayao fault zone while Tarlay earthquake initiated by Namma Fault Zone. To further refine seismic activities (b value) and seismic pattern (Dc) values including their relationship in this region, more detailed case studies of earthquakes that generated in different seismogenic faults are indispensable.

## 7. Acknowledgments

This research was supported by the National Research Council of Thailand (NRCT) 2017. I acknowledge thoughtful comments and suggestions by the editors and anonymous reviewers that enhanced the quality of this manuscript significantly.

## 8. References

- Aki, K., 1981. A probabilistic synthesis of precursory phenomena, in Earthquake Prediction: An International Review, D. W. Simpson and P. G. Richards (Editors), *American Geophysical Union, Washington, D.C.*; 566-574.
- Barton, D.J., Foulger, G.R., Henderson, J.R. and Julian, B.R., 1999. Frequency magnitude statistics and spatial correlation dimensions of earthquakes at Long Valley Caldera, California, *Geophys. J. Int.*; 138, 563-570.
- Bayrak, Y., Bayrak, E., 2012. Regional variations and correlations of Gutenberg-Richter parameters and fractal dimension for the different

- seismogenic zones in Western Anatolia, *Journal of Asian Earth Sciences.*; 58, 98–107.
- Bhattacharya, P., Kayal, J.R., Baruah, S. and Arefiev, S.S., 2010. Earthquake source zone in Northeast India: seismic tomography, fractal dimension and b value mapping, *Pure and Applied Geophysics.*; 167,999-1012.
- Chen, C., Wu, Y., 2006. An improved region-time-length algorithm applied to the 1999 Chi-Chi, Taiwan earthquake, *Geophys. J. Int.*; 166, 144-147.
- Chen, C., Wang, W., Chan, Y., Wu, Y. and Lee, Y., 2006. A correlation between the b-value and the fractal dimension from the aftershock sequence of the 1999 Chi-Chi, Taiwan, earthquake, *Geophys. J. Int.*; 167, 1215-1219.
- Fenton, C.H., Charusiri, P. and Wood, S.H., 2003. Recent paleoseismic investigations in Northern and Western Thailand. *Annals of Geophysics.*; 46(5): 957-981.
- Gardner, J. K. and Knopoff, L., 1974. Is the sequence of earthquakes in Southern California, with aftershocks removed, Poissonian?, *Bulletin Seismol. Soc. Am.*; 64(1), 363–367.
- Grassberger, P. and Procaccia, I., 1983. Measuring the strangeness of strange attractors, *Physica D.*; 9, 189-208.
- Gupta, I.D., 2002. The state of the art in seismic hazard analysis. *ISET J. Earthquake Technol.*, 39(428), 311–346.
- Gutenberg, B. and Richter, C.F., 1944. Frequency of earthquakes in California, *Bull Seismol Soc Am.*; 34, 185-188.
- Hirata, T., 1989. A correlation between the b-value and the fractal dimension of earthquakes, *J. geophys. Res.*, 94, 7507–7514.
- Kagan, Y.Y., Knopoff, L., 1980. Spatial distribution of earthquakes: the two-point correlation function, *Geophys. J. R. astr. Soc.*; 62, 303-320.
- Manakou, M. V. and Tsapanos, T. M., 2000. Seismicity and seismic hazard parameters evaluation in the island of Crete and surrounding area inferred from mixed files, *Tectonophysics.*; 321, 157-178.
- Mekong River Commission, 2010. State of the basin report 2010. *Mekong River Commission, Vientiane, Lao PDR.*
- Mogi, K., 1967. Earthquakes and fractures, *Tectonophysics.*; 5, 35-55.
- Molnar, P. and Topponnier, P., 1977. The collision between India and Eurasia, *Scientific American.*; 236, 4, 30-41.
- Nuannin, P., Kulhanek, O. and Persson, L., 2005. Spatial and temporal b value anomalies preceding the devastating off coast of NW Sumatra earthquake of December 26, 2004, *geophysical research letters.* ; 32, 11307.
- Pailoplee, S. and Choowong, M., 2014. Earthquake frequency-magnitude distribution and fractal dimension in mainland Southeast Asia, *Earth, Planets and space.*; 66,8.
- Pailoplee, S., Channarong, P. and Chutakositkanon, V., 2013. Earthquake activities in the Thailand-Laos-Myanmar border region: a statistical approach, *Terr. Atmos. Ocean. Sci.*; 24(4), Part II, 721-730.
- Pailoplee, S., Takashima, I., Kosuwan, S. and Charusiri, P., 2009. Earthquake activities along the Lampang-theon fault zone, Northern Thailand: evidence from paleoseismological and seismicity data, *Journal of Applied Sciences Research.*; 5(2), 168-180.
- Scholz, C.H., Aviles, C.A. and Wesnousky, S.G., 1986. Scaling differences between large interplate and intraplate

- earthquakes, *Bulletin of Seismological Society of America.*; 76(1), 65-70.
- Wiemer, S. and Wyss, M., 1994. Seismic quiescence before the Landers (M = 7.5) and Big Bear (M= 6.5) 1992 earthquakes. *Bulletin of the Seismological Society of America.*; 84, 900-916.
- Wu, Y., Chen, C. and Rundle, J.B., 2008. Precursory seismic activation of the Pingtung(Taiwan) offshore doublet earthquakes on 26 December 2006: A pattern informatics analysis, *Terr. Atmos. Ocean. Sci.*; 19, 6, 743-749.
- Woessner, J., Wiemer S., 2005. Assessing the Quality of Earthquake Catalogues: Estimating the Magnitude of Completeness and Its Uncertainty, *Bull. Seismol. Soc. Am.*; 95(2), 684-698.